

# The facial weak order in hyperplane arrangements

Aram Dermenjian<sup>1,3</sup>

Christophe Hohlweg<sup>1</sup>, Thomas McConville<sup>2</sup> and Vincent Pilaud<sup>3</sup>

<sup>1</sup>Université du Québec à Montréal (UQAM)

<sup>2</sup>Mathematical Sciences Research Institute (MSRI)

<sup>3</sup>École Polytechnique (LIX)

6 April 2019

On this day in 1896 the modern olympics began in Athens, Greece after being banned for over 1,500 years!

5

4

3

1

# The facial weak order in hyperplane arrangements

2

Aram Dermenjian<sup>1,3</sup>

Christophe Hohlweg<sup>1</sup>, Thomas McConville<sup>2</sup> and Vincent Pilaud<sup>3</sup>

<sup>1</sup>Université du Québec à Montréal (UQAM)

<sup>2</sup>Mathematical Sciences Research Institute (MSRI)

<sup>3</sup>École Polytechnique (LIX)


6 April 2019

On this day in 1896 the modern olympics began in Athens, Greece after being banned for over 1,500 years!

# History and Background - Hyperplanes

- $(V, \langle \cdot, \cdot \rangle)$  -  $n$ -dim real Euclidean vector space.
- A *hyperplane*  $H_i$  is codim 1 subspace of  $V$  with normal  $e_i$ .




## Example



# History and Background - Arrangements


- A *hyperplane arrangement* is  $\mathcal{A} = \{H_1, H_2, \dots, H_k\}$ .
- $\mathcal{A}$  is *central* if  $\{0\} \subseteq \bigcap \mathcal{A}$ .
- Central  $\mathcal{A}$  is *essential* if  $\{0\} = \bigcap \mathcal{A}$ .

## Example



## History and Background - Arrangements

- *Regions*  $\mathcal{R}$  - connected components of  $V$  without  $\mathcal{A}$ .
- *Faces*  $\mathcal{F}_{\mathcal{A}}$  - intersections of closures of some regions.




## History and Background - (Partial) Orders

- *Lattice* - poset where every two elements have a *meet* (greatest lower bound) and *join* (least upper bound).

### Example


The lattice  $(\mathbb{N}, |)$  where  $a \leq b \Leftrightarrow a | b$ .

- meet - greatest common divisor
- join - least common multiple




## History and Background - Poset of regions

- *Base region*  $B \in \mathcal{R}$  - some fixed region
- *Separation set for*  $R \in \mathcal{R}$   
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$



## History and Background - Poset of regions


- *Base region*  $B \in \mathcal{R}$  - some fixed region
- *Separation set for*  $R \in \mathcal{R}$   
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$





# History and Background - Poset of regions



- *Base region*  $B \in \mathcal{R}$  - some fixed region
- *Separation set for*  $R \in \mathcal{R}$   
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$
- *Poset of Regions*  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  where  
 $R \leq_{\mathcal{A}} R' \Leftrightarrow S(R) \subseteq S(R')$



## History and Background - Poset of regions

- A region  $R$  is *simplicial* if normal vectors for boundary hyperplanes are linearly independent.
- $\mathcal{A}$  is *simplicial* if all  $\mathcal{R}$  simplicial.

### Example






# History and Background - Poset of regions

Theorem (Björner, Edelman, Ziegler '90)

*If  $\mathcal{A}$  is simplicial then  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  is a lattice for any  $B \in \mathcal{R}$ . If  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  is a lattice then  $B$  is simplicial.*

## Example






# History and Background - Poset of regions

Theorem (Björner, Edelman, Ziegler '90)

*If  $\mathcal{A}$  is simplicial then  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  is a lattice for any  $B \in \mathcal{R}$ . If  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  is a lattice then  $B$  is simplicial.*

## Example



# Coxeter Arrangements

## Example

A *Coxeter arrangement* is the hyperplane arrangement associated to a Coxeter group.

<b>Coxeter Groups</b>		<b>Hyperplane Arrangements</b>
Reflecting hyperplanes	$\leftrightarrow$	Hyperplane arrangement
Root system	$\leftrightarrow$	Normals to hyperplanes
Inversion sets	$\leftrightarrow$	Separation sets
Weak order	$\leftrightarrow$	Poset of regions

# Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order of Coxeter groups to an order on all the faces of its associated arrangement for type  $A$ .
- In 2006, Palacios and Ronco extended this new order to Coxeter groups of all types using cover relations.
- In 2016, D, Hohlweg and Pilaud showed this extension has a global equivalent and produces a lattice in Coxeter arrangements.


# Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order of Coxeter groups to an order on all the faces of its associated arrangement for type  $A$ .
- In 2006, Palacios and Ronco extended this new order to Coxeter groups of all types using cover relations.
- In 2016, D, Hohlweg and Pilaud showed this extension has a global equivalent and produces a lattice in Coxeter arrangements.
- Questions: Can we extend this to hyperplane arrangements? Can we find both local and global definitions? When do we actually get a lattice?

# Facial Intervals

Proposition (Björner, Las Vergas, Sturmfels, White, Ziegler '93)

Let  $\mathcal{A}$  be central with base region  $B$ . For every  $F \in \mathcal{F}_{\mathcal{A}}$  there is a unique interval  $[m_F, M_F]$  in  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  such that

$$[m_F, M_F] = \{R \in \mathcal{R} \mid F \subseteq \overline{R}\}$$




# Facial Weak Order



Let  $\mathcal{A}$  be a central hyperplane arrangement and  $B$  a base region in  $\mathcal{R}$ .

## Definition



The *facial weak order* is the order  $\text{FW}(\mathcal{A}, B)$  on  $\mathcal{F}_{\mathcal{A}}$  where for  $F, G \in \mathcal{F}$ :

$$F \leq G \Leftrightarrow m_F \leq_{\mathcal{A}} m_G \text{ and } M_F \leq_{\mathcal{A}} M_G$$



# Facial Weak Order - Example





# Facial Weak Order - Example





# Facial Weak Order - Example





# Facial Weak Order - Example





# Facial Weak Order - Example



# Facial Weak Order - Example



# Facial Weak Order - Example







# Cover Relations

Proposition (D., Hohlweg, McConville, Pilaud, '18+)

For  $F, G \in \mathcal{F}_A$  if

1.  $F \leq G$  in  $\text{FW}(\mathcal{A}, B)$
2.  $|\dim(F) - \dim(G)| = 1$
3.  $F \subseteq G$  or  $G \subseteq F$

then  $F \lessdot G$ .




# Covectors

- *covector* - a vector in  $\{-, 0, +\}^A$  with signs relative to hyperplanes.
- $\mathcal{L} \subseteq \{-, 0, +\}^A$  - set of covectors

## Example

$$F_4 \leftrightarrow (+, 0, -) \quad F_4(H_1) = +; F_4(H_2) = 0; F_4(H_3) = -$$




# Covectors

- *covector* - a vector in  $\{-, 0, +\}^A$  with signs relative to hyperplanes.
- $\mathcal{L} \subseteq \{-, 0, +\}^A$  - set of covectors

## Example

$$F_4 \leftrightarrow (+, 0, -) \quad F_4(H_1) = +; F_4(H_2) = 0; F_4(H_3) = -$$





# Covector Definition

## Definition

For  $X, Y \in \mathcal{L}$ :

$$X \leq_{\mathcal{L}} Y \Leftrightarrow Y(H) \leq X(H) \quad \text{with } - < 0 < +$$



# Zonotopes

- *Zonotope*  $Z_{\mathcal{A}}$  is the convex polytope:

$$Z_{\mathcal{A}} := \left\{ v \in V \mid v = \sum_{i=1}^k \lambda_i e_i, \text{ such that } |\lambda_i| \leq 1 \text{ for all } i \right\}$$


## Theorem (Edelman '84, McMullen '71)

*There is a bijection between  $\mathcal{F}_{\mathcal{A}}$  and the nonempty faces of  $Z_{\mathcal{A}}$  given by the map*

$$\tau(F) = \left\{ v \in V \mid v = \sum_{F(H_i)=0} \lambda_i e_i + \sum_{F(H_j) \neq 0} \mu_j e_j \right\}$$

where  $|\lambda_i| \leq 1$  for all  $i$  and  $\mu_j = F(H_j)$

# Zonotope - Construction example





## Root inversion sets

■ roots  $\Phi_{\mathcal{A}} := \{\pm e_1, \pm e_2, \dots, \pm e_k\}$

■ root inversion set

$$\mathbf{R}(F) := \{e \in \Phi_{\mathcal{A}} \mid \langle x, e \rangle \leq 0 \text{ for some } x \in F\}.$$



# Equivalent definitions

## Theorem (D., Hohlweg, McConville, Pilaud '18+)

For  $F, G \in \mathcal{F}_{\mathcal{A}}$  the following are equivalent:





- $m_F \leq_{\mathcal{A}} m_G$  and  $M_F \leq_{\mathcal{A}} M_G$  in poset of regions  $(\mathcal{R}, B, \leq_{\mathcal{A}})$ .
- There exists a chain of covers in  $\text{FW}(\mathcal{A}, B)$  such that

$$F = F_1 \triangleleft F_2 \triangleleft \cdots \triangleleft F_n = G$$

- $F \leq_{\mathcal{L}} G$  in terms of covectors  $(G(H) \leq F(H) \forall H \in \mathcal{A})$
- $\mathbf{R}(F) \setminus \mathbf{R}(G) \subseteq \Phi_{\mathcal{A}}^-$  and  $\mathbf{R}(G) \setminus \mathbf{R}(F) \subseteq \Phi_{\mathcal{A}}^+$ .



# Equivalence for type $A_2$ Coxeter arrangement



## Facial weak order lattice


Theorem (D., Hohlweg, McConville, Pilaud '18+)

*The facial weak order  $\text{FW}(\mathcal{A}, B)$  is a lattice when  $\mathcal{A}$  is simplicial.*

Corollary (D., Hohlweg, McConville, Pilaud '18+)

*The lattice of regions is a sublattice of the facial weak order lattice when  $\mathcal{A}$  is simplicial.*

## Example: $B_3$ Coxeter arrangement




# Properties of the facial weak order

## Theorem (D., Hohlweg, McConville, Pilaud '18+)

- $\text{FW}(\mathcal{A}, B)$  is self-dual.
- $\mathcal{A}$  simplicial implies  $\text{FW}(\mathcal{A}, B)$  is semi-distributive.
- $\mathcal{A}$  simplicial and  $X \in \mathcal{F}_{\mathcal{A}}$  then  $X$  is join-irreducible in  $\text{FW}(\mathcal{A}, B)$  if and only if  $M_X$  is join-irreducible in  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  and  $\text{codim}(X) \in \{0, 1\}$
- Möbius function:  $X, Y \in \mathcal{F}_{\mathcal{A}}$  let  $Z = X \cap Y$ .

$$\mu(X, Y) = \begin{cases} (-1)^{\text{rk}(X) + \text{rk}(Y)} & \text{if } X \leq Z \leq Y \text{ and } Z = X_{-Z} \cap Y \\ 0 & \text{otherwise} \end{cases}$$



Thank you!